arXiv:1304.3281v1 [math-ph] 11 Apr 2013

ON PERIODIC WAVE FUNCTIONS OF SCHRÖDINGER OPERATORS ON CAYLEY TREES

FUMIO HIROSHIMA, UTKIR ROZIKOV

Abstract. In the paper we define periodic wave functions for a (discrete) Schrödinger operator on a Cayley tree. This periodicity depends on a subgroup of a group representation of the Cayley tree. For any subgroup of finite index we give a criterion for eigenvalues of the Schrödinger operator under which there are periodic wave functions. For a normal subgroup of infinite index we describe a class of periodic wave functions.

Mathematics Subject Classifications (2010). 82B26.
Key words. Cayley tree, Anderson model, Schrödinger equation, periodic wave function.

1. Introduction

The topics of statistical mechanics on non-amenable graphs is growing (see for example, [11]). One of the non-amenable graph is a Cayley tree. The Cayley tree is not a realistic lattice, however, its amazing topology makes the exact calculation of various quantities possible [11]. It is believed that several among its interesting thermal properties could persist for regular lattices, for which the exact calculation is far intractable.

The spectral properties of Schrödenger operators on graphs have many applications in physics and they have been intensively studied since late 1990s. We shall consider Schrödinger operators which correspond to the Anderson model on a Cayley tree. In [9] Klein proved the existence of purely absolutely continuous spectrum, under weak disorder, on the Cayley tree. A new proof of a version of Kleins theorem is given in [2].

The paper [7] reviews the main aspects and problems in the Anderson model on the Cayley tree. It shows how the question as to whether wave functions are extended or localized is related to the existence of complex solutions of a certain nonlinear equation. In [10] sufficient conditions for the presence of the absolutely continuous spectrum of a Schrödinger operator on a rooted Cayley tree are given. See [1] for more results on the model. In [6] the spectrum of Schrödinger operator on the d-dimensional lattice $\mathbb{Z}^d$ is studied.

In this paper we construct some periodic wave functions for the Schrödinger operators which correspond to the Anderson model on a Cayley tree. The paper is organized as follows. In Section 2 we give some preliminaries (definition of Cayley tree, a group representation of a Cayley tree of order $k \geq 1$, its partition with respect to an arbitrary subgroup, the model). In Section 3 we define periodic wave functions. This periodicity depends on a subgroup of a group representation of the Cayley tree. For any subgroup of finite index we give a criterion for eigenvalues of the Schrödinger operator under which



FIGURE 1. Cayley tree of order two, i.e., $k = 2$.

there are periodic wave functions. For a normal subgroup of infinite index we describe a class of periodic wave functions.

2. PRELIMINARIES

Cayley tree. The Cayley tree Γ^k of order $k \geq 1$ is an infinite tree, i.e., a graph without cycles, such that exactly $k+1$ edges originate from each vertex (see Fig. 1). Let $\Gamma^k = (V, L)$ where V is the set of vertices and L the set of edges. Two vertices x and y are called *nearest neighbors* if there exists an edge $l \in L$ connecting them. We will use the notation $l = \langle x, y\rangle$.

Denote $S(x) = \{y \in V : \langle x, y\rangle\}$ the set of all neighbors of x.

A group representation of the Cayley tree.

Let G_k be a free product of $k+1$ cyclic groups of the second order with generators $a_1, a_2, \ldots, a_{k+1}$, respectively. Any element $x \in G_k$ has the following form:

$$x = a_{i_1}a_{i_2}\ldots a_{i_n}, \quad \text{where} \ \ 1 \leq i_m \leq k+1, \, m = 1, \ldots, n.$$

It is known that there exists a one-to-one correspondence between the set of vertices V of the Cayley tree Γ^k and the group G_k. In Fig. 1 some elements of G_k are shown.

In the group G_k, let us consider the left (right) shift transformations defined as follows. For $g_0 \in G_k$, let us set

$$T_g(h) = gh, \quad (T_g(h) = hg), \quad \text{for all} \ \ h \in G_k. \tag{2.1}$$

The set of all left (right) shifts in G_k is isomorphic to the group G_k.

Normal subgroups of finite index. In this subsection we answer to the following question: Does G_k have subgroups (in particular, normal subgroups) of any finite index?

Let $N_k = \{1, \dots, k+1\}$.

For any $\emptyset \neq A \subseteq N_k$, there exists a subgroup $H_A \subset G_k$ with the following properties:

(a) H_A is a normal subgroup with index $|G_k : H_A| = 2$;
(b) $H_A \neq H_B$ for all $A \neq B \subseteq N_k$.

Denote by e the unity element of G_k. Let $e \neq x_0 \in G_k$. Then there exists a normal finite-index subgroup H_{x_0}, which does not contain the element x_0.

The following hold:

1. If x is generated by more then one generators, then $|G_k : H_x| \geq 4$;
2. $|G_k : H_{a_i a_j}| = 6$, for any $i \neq j \in N_k$.
3. The group G_k does not have normal subgroups of odd index ($\neq 1$).
4. The group G_k has a normal subgroups of arbitrary even index.

Subgroups of infinite index. There are normal subgroups of infinite index. Some of them can be described as follows. Fix $M \subseteq N_k$ such that $|M| > 1$. $|\bullet|$ is the cardinality of $\bullet$. Let the mapping $\pi_M : \{a_1, ..., a_{k+1}\} \longrightarrow \{a_i,\ i \in M\} \cup \{e\}$ be defined by

$$\pi_M(a_i) = \begin{cases} a_i, & \text{if } \ i \in M \\ e, & \text{if } \ i \notin M. \end{cases}$$

Denote by G_M the free product of cyclic groups $\{e, a_i\},\ i \in M$. Consider

$$f_M(x) = f_M(a_{i_1} a_{i_2} ... a_{i_m}) = \pi_M(a_{i_1}) \pi_M(a_{i_2}) ... \pi_M(a_{i_m}).$$

Then it is easy to see that f_M is a homomorphism and hence $H_M = \{x \in G_k :\ f_M(x) = e\}$ is a normal subgroup of infinite index.

Partition structures of the Cayley tree.

Let $H_0 \subset G_k$ be an arbitrary subgroup of index n of the group G_k. Obviously, each subgroup of the group G_k is the kernel of some homomorphism φ of the group G_k into some group G^*. Introduce the following equivalence relation on the set G_k: $x \sim y$ if $xy^{-1} \in H_0$.

Denote by $S(x) = \{y \in G_k : \langle x, y \rangle\}$ the set of all nearest neighbors of the word $x \in G_k$. Let $G_k / H_0 = \{H_0, H_1, \dots, H_{n-1}\}$ be the factor-group w.r.t. H_0. In addition, let $q_i(x) = |S(x) \cap H_i|$, $i = 0, \dots, n-1$, and $Q(x) = (q_0(x), q_1(x), \dots, q_{n-1}(x))$, $x \in G_k$.

If $x \sim y$, then $q_i(x) = q_i(y)$ for $i = 0, \dots, n-1$, thus $Q(x) = Q(y)$.

Introduce the following notations: $q_i(H_0) = q_i(e) = |\{j : a_j \in H_i\}|$, $Q(H_0) = (q_0(H_0), \dots, q_{n-1}(H_0))$ and $N(H_0) = |\{j : q_j(H_0) \neq 0\}|$.

For any $x \in G_k$, there exists a permutation π_x of the coordinates of the vector $Q(H_0)$ such that

$$\pi_x Q(H_0) = Q(x). \tag{2.2}$$

Let $N(x) = |\{j : q_j(x) \neq 0\}|$. For any $x \in G_k$ we have $N(x) = N(H_0)$.

Remark. The group representation was given by Ganikhodjaev [3]. In [4] the classes of finite-index subgroups of the group G_k were constructed. In [8] the assertion that the group G_k does not have normal subgroups of odd index ($\neq 1$) is proved. Examples of normal subgroups of infinite index are constructed in [13]. In [12] the structure of partitioning of the group G_k into adjacent classes w.r.t. a normal subgroup of finite index was described.

Anderson model.

The Anderson Hamiltonian, H, on the Hilbert space

$$\ell^2(\Gamma^k)=\{\varphi:V\to\mathbb{C}:\sum_{x\in V}|\varphi(x)|^2<\infty\}$$

is the operator of the form $H=\Delta+\epsilon v$, where

1. The free Laplacian Δ is defined by

$$(\Delta\varphi)(x)=\sum_{y:\,d(x,y)=1}(\varphi(x)-\varphi(y)),\quad \text{for all}\quad \varphi\in\ell^2(\Gamma^k),$$

   with the usual distance d denoting the number of edges in the shortest (only) path between sites on the Cayley tree.

2. The operator v is a random potential,

$$(v\varphi)(x)=v(x)\varphi(x),$$

   where $v=\{v(x)\}_{x\in V}$ is a family of independent, identically distributed real random variables with common probability distribution ν. The coupling constant ϵ measures the disorder.

The goal is to study spectrum for this operator. The Schrödinger equation reads

$$\sum_{y\in S(x)}\varphi(y)=E\varphi(x)-\epsilon v(x)\varphi(x). \tag{2.3}$$

Here $\varphi(x)$ is the value of the wave function at vertex x, E is the energy of the particle (eigenvalue of the operator H), and the sum is over the $k+1$ neighbors of the site x.

3. PERIODIC WAVE FUNCTIONS.

Since the set of vertices V has the group representation G_k. Without lost of generality we identify V with G_k, i.e., we sometimes replace V with G_k.

Now give some periodic solutions to (2.3).

Definition 1. *Let K be a subgroup of $G_k, k\geq 1$. We say that a functions $\varphi=\{\varphi(x)\in\mathbb{C}:x\in G_k\}$ is* K-periodic *if $\varphi(yx)=\varphi(x)$ for all $x\in G_k$ and $y\in K$.*

A G_k-periodic function φ is called translation-invariant.

3.1. **The case of finite index.**

Let $G_k/K=\{K_1,...,K_r\}$ be a factor group, where K is a normal subgroup of index $r\geq 1$.

Note that any K-periodic function φ has the form

$$\varphi(x)=\varphi_j,\quad \forall x\in K_j,\quad j=1,\dots,r. \tag{3.1}$$

We note that the functions $q_j(x)$ introduced in the previous section are periodic, i.e. $q_j(x)=q_{ij}$, for all $x\in K_i$. Assume that $v=\{v(x)\}_{x\in V}$ is K-periodic, i.e. $v(x)=v_j$ for any $x \in K_j$ then from (2.3) we get

$$\sum_{j=1}^{r} q_{ij}\varphi_j = (E - \epsilon v_i)\varphi_i, \quad i = 1, 2, \dots, r. \tag{3.2}$$

This system can be written as

$$\sum_{\substack{j=1 \\ j \neq i}}^{r} q_{ij}\varphi_j - (E - q_{ii} - \epsilon v_i)\varphi_i = 0, \quad i = 1, 2, \dots, r. \tag{3.3}$$

For a given normal subgroup K of index $r \geq 1$ we denote by $D_K(E, v, \epsilon)$ the determinant of the system (3.3).

Thus the following is true

Theorem 1. *For given K, ϵ and a K-periodic v there exists a K-periodic wave function $\varphi \neq 0$ if and only if E is a solution to $D_K(E, v, \epsilon) = 0$.*

Examples. 1) In case $K = G_k$ we have translation invariant wave function, i.e. $\varphi(x) = \varphi_1$ for all $x \in V$. In this case $D_K(E, v, \epsilon) = E - \epsilon v_1 - k - 1 = 0$. Thus translation invariant wave function exists if and only if $E = \epsilon v_1 + k + 1$.

2) Let $K = G_k^{(2)}$ be the subgroup in G_k consisting of all words of even length. Clearly, $G_k^{(2)}$ is a subgroup of index 2. In this case we have $D_K(E, v, \epsilon) = (E - \epsilon v_1)(E - \epsilon v_2) - (k+1)^2 = 0$. For any E satisfying this equation there are $G_k^{(2)}$-periodic wave functions having the form

$$\varphi(x) = \begin{cases} \varphi_1, & \text{if } x \in G_k^{(2)} \\ \varphi_2, & \text{if } x \in G_k \setminus G_k^{(2)}. \end{cases}$$

3) Let $k = 2$, $K = H_0$, with $H_0 = H_{\{1\}} \cap H_{\{2\}} = \{x \in G_2 : \omega_x(a_1) - \text{even}, \ \omega_x(a_2) - \text{even}\}$, where the number of letters a_i of the word x is denoted by $\omega_x(a_i)$.

Note that H_0 is a normal subgroup of index 4. The group $G_2/H_0 = \{H_0, H_1, H_2, H_3\}$ has the following elements:

$$H_1 = \{x \in G_2 : \omega_x(a_1) - \text{even}, \ \omega_x(a_2) - \text{odd}\},$$
$$H_2 = \{x \in G_2 : \omega_x(a_1) - \text{odd}, \ \omega_x(a_2) - \text{even}\},$$
$$H_3 = \{x \in G_2 : \omega_x(a_1) - \text{odd}, \ \omega_x(a_2) - \text{odd}\}.$$

In the Fig. 2, the partitions of Γ^2 with respect to $H_0 = H_{\{1\}} \cap H_{\{2\}}$ is given. The elements of the class H_i, $i = 0, 1, 2, 3$ denoted by i.

In this case H_0-periodic wave function φ has the form $\varphi(x) = \varphi_i$ if $x \in H_i$. Such a function exists iff E satisfies the following equation

$$D_{H_0}(E, v, \epsilon) = \begin{vmatrix} \epsilon v_0 + 1 - E & 1 & 1 & 0 \\ 1 & \epsilon v_1 + 1 - E & 0 & 1 \\ 1 & 0 & \epsilon v_2 + 1 - E & 1 \\ 0 & 1 & 1 & \epsilon v_3 + 1 - E \end{vmatrix} = 0.$$



FIGURE 2. Partition of Γ^2 by subgroup H_0 of index 4.

3.2. **The case of infinite index.** We consider a normal subgroup $\mathcal{H}_0$ of infinite index constructed as in previous section with $M=\{1,2\}$. The factor group has the following form

$$G_k/\mathcal{H}_0=\{\mathcal{H}_0,\mathcal{H}_0(a_1),\mathcal{H}_0(a_2),\mathcal{H}_0(a_1a_2),\dots\},$$

where $\mathcal{H}_0(y)=\{x\in G_k: f_M(x)=y\}$. We introduce the notations

$$\mathcal{H}_n=\mathcal{H}_0(\underbrace{a_1a_2\dots}_{n}),\quad \mathcal{H}_{-n}=\mathcal{H}_0(\underbrace{a_2a_1\dots}_{n}).$$

In this notation, the factor group can be represented as

$$G_k/\mathcal{H}_0=\{\dots,\mathcal{H}_{-2},\mathcal{H}_{-1},\mathcal{H}_0,\mathcal{H}_1,\mathcal{H}_2,\dots\}.$$

The partition of the Cayley tree Γ^2 w.r.t. $\mathcal{H}_0$ is shown in Fig. 3 (the elements of the class $\mathcal{H}_i$, $i\in\mathbb{Z}$, are merely denoted by i).

Any $\mathcal{H}_0$-periodic wave function has the form:

$$\varphi(x)=\varphi_n \ \text{ if } \ x\in\mathcal{H}_n,\quad n\in\mathbb{Z}.$$

We note that (see [14]) if $x\in\mathcal{H}_n$ then $S(x)\subset\mathcal{H}_{n-1}\cup\mathcal{H}_n\cup\mathcal{H}_{n+1}$. Moreover

$$|S(x)\cap\mathcal{H}_{n-1}|=1,\quad |S(x)\cap\mathcal{H}_n|=k-1,\quad |S(x)\cap\mathcal{H}_{n+1}|=1.$$



FIGURE 3. The partition of the Cayley tree Γ^2 w.r.t. $\mathcal{H}_0$, the elements of the class $\mathcal{H}_i$, $i \in \mathbb{Z}$, are denoted by i.

Assume that v is $\mathcal{H}_0$-periodic, i.e. $v(x) = v_n$ for any $x \in \mathcal{H}_n$. Then from equation (2.3) we get (see Fig. 3)

$$(E - \epsilon v_n)\varphi_n = \varphi_{n-1} + (k-1)\varphi_n + \varphi_{n+1}, \tag{3.4}$$

where $n \in \mathbb{Z}$.

For simplicity we assume that $v_n \equiv v_0$, i.e. v is translation invariant. Then equation (3.4) has the following general solution

$$\varphi_n = C_1\lambda_1^n + C_2\lambda_2^n,$$

where λ_1, λ_2 are solution to equation

$$\lambda^2 + (k - 1 + \epsilon v_0 - E)\lambda + 1 = 0.$$

Thus we have proved the following

Theorem 2. *For $\mathcal{H}_0$, given ϵ and translation invariant v there exists a $\mathcal{H}_0$-periodic wave function φ, such that $\varphi(x) = C_1\lambda_1^n + C_2\lambda_2^n$ for any $x \in \mathcal{H}_n$, $n \in \mathbb{Z}$.*

REFERENCES

[1] B. Derrida, G.J. Rodgers, *Anderson model on a Cayley tree: the density of states.* J. Phys. A **26**(9) (1993), L457–L463.

[2] R. Froese, D. Hasler, W. Spitzer, *Absolutely continuous spectrum for the Anderson model on a tree: a geometric proof of Klein's theorem.* Comm. Math. Phys. **269**(1) (2007), 239–257.

[3] N. N. Ganikhodjaev, *Group presentations and automorphisms of the Cayley tree*, Doklady AN Uzbekistan., 4 (1994), 3–5.

[4] N.N. Ganikhodjaev, U.A. Rozikov, *Classes of normal subgroups of finite index of the group representation of the Cayley tree.* Uzbek. Mat. Jour., 4 (1997), 31–39.(Russian)

[5] N.N. Ganikhodjaev, U.A. Rozikov, *A description of periodic extremal Gibbs measures of some lattice models on the Cayley tree*, Theor. Math. Phys. **111** (1997), 480–486.

[6] F.Hiroshima, I.Sasaki, T.Shirai, A.Suzuki, *Note on the spectrum of discrete Schrödinger operators.* 2012, arXiv:1209.0522.

[7] J. Miller, B. Derrida, *Weak-disorder expansion for the Anderson model on a tree.* J. Statist. Phys. **75**(3-4) (1994), 357–388.

[8] E.P. Normatov, U.A. Rozikov, *Description of harmonic functions using group representations of the Cayley tree.* Math. Notes., **79**(3-4) (2006), 399–407.

[9] A. Klein. *Extended states in the Anderson model on the Bethe lattice.* Advances in Math., **133** (1998), 163–184.

[10] S.Kupin, *Absolutely continuous spectrum of a Schrödinger operator on a tree.* J. Math. Phys., **49** (2008), 113506-10.

[11] R.Lyons, *Phase transitions on nonamenable graphs.* J. Math. Phys. **41** (2000), 1099–1126.

[12] U.A. Rozikov, *Structures of partitions of the group representation of the Cayley tree into cosets by finite-index normal subgroups, and their applications to the description of periodic Gibbs distributions*, Theor. Math. Phys. **112**(1) (1997), 929–933.

[13] U.A. Rozikov, *Countably periodic Gibbs measures of the Ising model on the Cayley tree.* Theor. Math. Phys. **130**(1) (2002), 92–100.

[14] U.A. Rozikov, F.T. Ishankulov, *Description of periodic p-harmonic functions on Cayley trees.* NoDEA. Nonlinear Differential Equations and Appl. **17**(2) (2010), 153–160.

F. Hiroshima, Faculty of Mathematics, Kyushu University, Fukuoka, 819-0395, Japan.
E-mail address: `hiroshima@math.kyushu-u.ac.jp`

U. A. Rozikov, Institute of mathematics, 29, Do'rmon Yo'li str., 100125, Tashkent, Uzbekistan.
E-mail address: `rozikovu@yandex.ru`